arXiv:0712.0664v1 [hep-ph] 5 Dec 2007

Frascati Physics Series Vol. XLVI (2007), pp. 000-000
HADRON07: XII Int. Conf. on Hadron Spectroscopy – Frascati, October 8-13, 2007
Parallel Session

# ON THE EXISTENCE OF LIGHT-SCALAR MESONS $\kappa(800)$ and $\kappa'(1150)$: THE $\widetilde{U}(12)$ SCHEME AND BES II DATA

Kenji Yamada and Tomohito Maeda
*Department of Engineering Science, Junior College Funabashi Campus,*
*Nihon University, Funabashi 274-8501, Japan*

## Abstract

We present that there should exist a light strange-scalar meson $\kappa'$, in addition to the $\kappa(800)$, which has a mass around 1.1-1.2 GeV, a rather narrow width, and couples strongly to $\kappa(800)\sigma(600)$ $(K\pi\pi\pi)$ but weakly to $K\pi$, based upon the $\widetilde{U}(12)$-classification scheme of hadrons and BES II data on $J/\psi \to \bar{K}^*(892)^0 K^+\pi^-$ decay.

## 1 Introduction

Recently, the existence of the light-scalar mesons, $\sigma(600)$ and $\kappa(800)$, has been confirmed by showing the presence of respective poles in the $\pi\pi$ 1, 2) and $K\pi$ 2, 3) scattering amplitudes, in addition to results of Breit-Wigner fits to $D$- and $J/\psi$-decay data, respectively, from the E791 4) and BES 5, 6) collaborations. However, the nature of these resonances, together with the

$f_0(980)$ and $a_0(980)$, has been a long-standing problem in controversy, where it is not obvious how these light-scalar mesons are understood in terms of quarks and gluons in QCD.

Here we focus on the strange scalar mesons and discuss the existence of an extra $\kappa'$ meson, in addition to the normal $\kappa(800)$, and their strong decay properties.

## 2 Existence of the extra $\kappa'$ meson

### 2.1 The $\widetilde{U}(12)$-classification scheme of hadrons

The $\widetilde{U}(12)$-classification scheme of hadrons, 7, 8) which has a manifestly covariant framework of $\widetilde{U}(12)_{SF} \times O(3,1)_L$, generalized covariantly from nonrelativistic $SU(6)_{SF} \times O(3)_L$ by boosts, separating the spin and space degrees of freedom, gives covariant quark representations for composite hadrons with definite Lorentz and chiral transformation properties. The $\widetilde{U}(12)$-classification scheme has a "static" unitary $U(12)_{SF}$ spin-flavor symmetry in the rest frame of hadrons, embedded in the covariant $\widetilde{U}(12)$-representation space, where $\widetilde{U}(12)$ has as its subgroups the pseudounitary homogeneous Lorentz group for Dirac spinors and unitary symmetry group for light-quark flavors,

$$\widetilde{U}(12)_{SF} \supset \widetilde{U}(4)_D \times U(3)_F. \tag{1}$$

Since

$$U(12)_{SF} \supset U(4)_D \times U(3)_F, \tag{2a}$$

$$U(4)_D \supset SU(2)_\rho \times SU(2)_\sigma, \tag{2b}$$

the static $U(12)_{SF}$ symmetry includes both the nonrelativistic spin-flavor $SU(6)_{SF}$ and chiral $U(3)_L \times U(3)_R$ symmetry[1] as

$$U(12)_{SF} \supset SU(6)_{SF} \times SU(2)_\rho, \tag{3a}$$

$$U(12)_{SF} \supset U(3)_L \times U(3)_R \times SU(2)_\sigma, \tag{3b}$$

---

[1] Hadron states are classified, aside from flavors, by the quantum numbers $\rho, S, L, J, P$, where $\rho$ is the net quark $\rho$-spin concerning $SU(2)_\rho$, $S$ the ordinary net quark $\sigma$-spin, $L$ the total quark orbital angular momentum, and $J$ and $P$ the total spin and parity of hadrons.

where $SU(2)_\rho$ and $SU(2)_\sigma$ are the Pauli-spin groups concerning the boosting and intrinsic spin rotation, respectively, of constituent quarks, being connected with decomposition of Dirac $\gamma$-matrices, $\gamma = \rho \times \sigma$. This implies that the $\widetilde{U}(12)$-classification scheme is able to incorporate effectively, according to dynamical consequences of QCD, the effects of chiral symmetry and its spontaneous breaking, essential for understanding of properties of the low-lying hadrons, into what is called a constituent quark model.

In the $\widetilde{U}(12)$-classification scheme there are two light-scalar meson multiplets, normal $S^{(N)}$ and extra $S^{(E)}$ with $J^{PC} = 0^{++}$ and $0^{+-}$, respectively, in the ground level ($L = 0$). These $N$- and $E$-scalar multiplets are the chiral partners, respectively, of the $N$- and $E$-pseudoscalar multiplets and they form linear representations of the $U(3)_L \times U(3)_R$ chiral symmetry. Concerning the strange scalar mesons, now we have two $\kappa$ mesons, $\kappa^{(N)}(0^{++})$ and $\kappa^{(E)}(0^{+-})$. Note that the observed $\kappa(800)$ and missing $\kappa'$ are generally mixtures of them. 9)

### 2.2 The BES II data

The $K^+\pi^-$ mass spectrum in $J/\psi \to \bar{K}^*(892)^0 K^+\pi^-$ decay observed by the BES II experiment 6) is shown in fig.1 where there seems to be a visible bump structure at 1.1-1.2 GeV. If this structure is attributed to the existence of a new $K\pi$ resonance, its spin-parity will likely be $0^+$ or $1^-$, since higher spins are unfavorable for such a low-mass state, and also its width is supposed to be narrow, judging from the data structure.

We hereafter refer to the strange scalar meson mentioned above as the $\kappa'(1150)$.[2]

## 3 Strong decays of the $\kappa(800)$ and $\kappa'(1150)$ mesons

We examine strong two-body decays of the $\kappa(800)$ and $\kappa'(1150)$ as mixtures of the $\kappa^{(N)}$ and $\kappa^{(E)}$ in the $\widetilde{U}(12)$-classification scheme as follows:

$$\kappa(800) \to K + \pi \tag{4}$$

---

[2] A recent lattice-QCD study on light-scalar mesons by the UKQCD collaboration 10) suggests that the $a_0(980)$ is predominantly a conventional $\bar{q}q$ state, while the $\kappa(800)$ is too light to be assigned to the $\bar{q}q$ state, which is expected to have a mass about 100-130 MeV heavier than the $a_0(980)$.



Figure 1: *The* $K^+\pi^-$ *invariant mass spectrum in* $J/\psi \to \bar{K}^*(892)^0 K^+\pi^-$ *decay from BES II.* $^{6)}$ *The dark shaded histograms show contribution from the* $\kappa(800)$.

and

$$\kappa'(1150) \to K + \pi, \tag{5a}$$

$$\to K + \eta, \tag{5b}$$

$$\to \kappa(800) + \sigma(600) \quad [\to K\pi\pi\pi]. \tag{5c}$$

In the actual calculations of decay matrix elements we treat the strange mesons $K$, $\kappa$ and $\kappa'$ as quark-composite $n\bar{s}$ states, while $\pi$, $\eta$ and $\sigma$ as external local fields.

### 3.1 Quark-pseudoscalar and quark-scalar couplings

For the effective quark-pseudoscalar coupling inside hadrons we assume the two independent interactions of the forms

$$g_{ps}\bar{q}(-i\gamma_5)q\phi_p \qquad \text{for pseudoscalar type,} \tag{6a}$$

$$g_{pv}\bar{q}(-i\gamma_5\gamma_\mu)q\partial_\mu\phi_p \qquad \text{for pseudovector type.} \tag{6b}$$

The effective quark-scalar coupling is simply related to the quark-pseudoscalar coupling, assuming the $\sigma$ meson is a chiral partner of the $\pi$ meson in the linear representation of chiral symmetry,[3] and given as

$$g_{ps}\bar{q}q\phi_\sigma, \tag{7a}$$
$$g_{pv}\bar{q}\gamma_\mu q\partial_\mu\phi_\sigma. \tag{7b}$$

Then the matrix elements for the pseudoscalar($\pi, \eta$)-emitted processes are generally given by

$$T^{(P)} = T^{(P)}_{ps} + T^{(P)}_{pv} \tag{8a}$$

with

$$T^{(P)}_{ps} = g_{ps}\langle \overline{W}(v')(-i\gamma_5\phi_p)W(v)iv\cdot\gamma\rangle I^{(P)}_G(q^2) + \text{c.c.}, \tag{8b}$$
$$T^{(P)}_{pv} = g_{pv}\langle \overline{W}(v')(-\gamma_5\gamma_\mu q_\mu\phi_p)W(v)iv\cdot\gamma\rangle I^{(P)}_G(q^2) + \text{c.c.}, \tag{8c}$$

where $W(v)$ and $\overline{W}(v')$ are the spin wave functions of initial- and final-state mesons, 7) $I^{(P)}_G(q^2)$ a space-time part of the Lorentz-invariant transition form factor,[4] $v$ and $v'$ the 4-velocities, $q_\mu$ the momentum of emitted pseudoscalar mesons, and $\langle\cdots\rangle$ means the trace taken over the spinor and flavor indices.

The matrix elements for the $\sigma$-emitted processes are given likewise by

$$T^{(\sigma)} = T^{(\sigma)}_{ps} + T^{(\sigma)}_{pv} \tag{9a}$$

with

$$T^{(\sigma)}_{ps} = g_{ps}\langle \overline{W}(v')(\phi_\sigma)W(v)iv\cdot\gamma\rangle I^{(\sigma)}_G(q^2) + \text{c.c.}, \tag{9b}$$
$$T^{(\sigma)}_{pv} = g_{pv}\langle \overline{W}(v')(-i\gamma_\mu q_\mu\phi_\sigma)W(v)iv\cdot\gamma\rangle I^{(\sigma)}_G(q^2) + \text{c.c.}. \tag{9c}$$

### 3.2 Evaluation of the coupling constants

The coupling constants $g_{ps}$ and $g_{pv}$ were evaluated by Maeda et al.. 11) They calculated the $D$-wave/$S$-wave amplitude ratio and width of $b_1(1235) \rightarrow \omega(782) + \pi$ decay and obtained the values $g_{ps} = 2.07$ GeV and $g_{pv} = 14.0$, using their experimental values $D/S = 0.277$ ($\pm 0.027$) and $\Gamma[\omega\pi] \approx \Gamma^{\text{tot}} = 142$ ($\pm 9$) MeV 12) as input.

---

[3] Here we take chiral $SU(2)_L \times SU(2)_R$ as imposed symmetry.

[4] This is obtained from overlap integral of space-time wave functions for initial and final mesons. In the present analysis we simply take $I^{(P)}_G = I^{(\sigma)}_G = 1$, since initial and final mesons both belong to the ground-state ($L = 0$) multiplet in the $\widetilde{U}(12)_{SF} \times O(3,1)_L$-classification scheme.

We now calculate the decay width of $K^*(892) \to K+\pi$ to see the validity of the present decay model and obtain a reasonable value of $\Gamma[K\pi] = 58$ MeV, compared with the experimental value $\Gamma[K\pi] \approx \Gamma^{\rm tot} = 50.8 \pm 0.9$ MeV. 12)

### 3.3 Strong decay widths of the $\kappa(800)$ and $\kappa'(1150)$

Since the $\kappa(800)$ and $\kappa'(1150)$ are generally mixtures of $\kappa^{(N)}$ and $\kappa^{(E)}$, we introduce the mixing angle $\theta$, which is the only free parameter in the present analysis, by

$$|\kappa(800)\rangle = \cos\theta\ |\kappa^{(E)}\rangle + \sin\theta\ |\kappa^{(N)}\rangle, \tag{10a}$$

$$|\kappa'(1150)\rangle = -\sin\theta\ |\kappa^{(E)}\rangle + \cos\theta\ |\kappa^{(N)}\rangle. \tag{10b}$$

Here we take the mixing angle $\theta$ to be around $-65^\circ$ so that the $\kappa(800)$ has a width of several hundred MeV and the $\kappa'(1150)$ a rather narrow width, in conformity with their observed properties, $\Gamma[\kappa(800)] = 550 \pm 34$ MeV 12) and the BES II data mentioned above for the $\kappa'(1150)$.

Using the mixing angle $\theta = -65^\circ$, we evaluate the partial decay widths of the $\kappa(800)$ and $\kappa'(1150)$ for respective channels in the following.

#### *3.3.1 Decay of the* $\kappa(800)$

If we take a mass of the $\kappa(800)$ to be 800 MeV, we obtain $\Gamma[K\pi] = 354$ MeV for $\kappa(800) \to K+\pi$. This is consistent, though somewhat small, with the experimental value $\Gamma^{\rm tot} \approx \Gamma[K\pi] = 550 \pm 34$ MeV.

#### *3.3.2 Decays of the* $\kappa'(1150)$

We take tentatively 1150 MeV for a mass of the missing state $\kappa'(1150)$ and then obtain

$$\Gamma[K\pi] = 18 \text{ MeV} \quad \text{for} \quad \kappa'(1150) \to K+\pi, \tag{11a}$$

$$\Gamma[K\eta] = 2 \text{ MeV} \quad \text{for} \quad \kappa'(1150) \to K+\eta, \tag{11b}$$

$$\Gamma[\kappa\sigma] = 30 \text{ MeV} \quad \text{for} \quad \kappa'(1150) \to \kappa(800)+\sigma(600), \tag{11c}$$

where mass values of the $\kappa(800)$ and $\sigma(600)$ are taken tentatively to be 600 MeV and 350 MeV, respectively, and the singlet-octet mixing angle for the pseudoscalar mesons $\eta$ and $\eta'$ to be $\theta_P = -11.5^\circ$. From these partial decay widths we could estimate the total width to be

$$\Gamma^{\rm tot} \approx \Gamma[K\pi] + \Gamma[K\eta] + \Gamma[\kappa\sigma] = 50 \text{ MeV}. \tag{12}$$

It may be worthwhile to mention that the dominant decay mode of the $\kappa'(1150)$ is not $K\pi$ but $\kappa(800)\sigma(600)$ ($K\pi\pi\pi$), the $K\pi$ branching ratio is $\Gamma[K\pi]/\Gamma^{\rm tot} \approx 0.36$, and therefore the $\kappa'(1150)$ is supposed not to be seen in the $K\pi$ scattering processes.[5]

Here it goes without saying that the present treatment of the $\kappa(800)$ and $\sigma(600)$ as narrow resonances is quite a rough approximation and the evaluated decay width to $\kappa(800)\sigma(600)$ does not really make sense. In practice we should perform a dynamical calculation of the decay chain $\kappa'(1150) \to \kappa(800)\sigma(600) \to K\pi\pi\pi$, taking into account the effects of the broad $\kappa$ and $\sigma$ widths.

## 4 Concluding Remarks

We presented that there should exist an extra $\kappa'$ meson which has a mass around 1.1-1.2 GeV, a rather narrow width, and couples strongly to $\kappa(800)\sigma(600)$ ($K\pi\pi\pi$) but weakly to $K\pi$, based upon the $\widetilde{U}(12)$-classification scheme and BES II data.

The strong coupling to $\kappa(800)\sigma(600)$ suggests that to observe the $\kappa'$ meson experimentally it might be favorable to study the $K\pi\pi\pi$ system, for example, in $J/\psi \to K^*(892)(K\pi\pi\pi)$ decay and $e^+e^- \to K^*(892)(K\pi\pi\pi)$ processes. However, if the main component of $\kappa'$ is $\kappa^{(E)}(0^{+-})$, as is in the present analysis (taking $\theta = -65^\circ$ in eq.10b), the $\kappa'$ production in these processes[6] would be suppressed by charge-conjugation ($C$) invariance in the limit of $SU(3)_f$ symmetry. 14) Rather, the $\chi_{c0,1,2} \to K^*(892)(K\pi\pi\pi)$ decay processes would be more promising, since they are $C$-parity allowed decays in the $SU(3)_f$ limit.

---

[5] This is consistent with experimental data 13) on the $S$-wave phase of the $K\pi$ scattering amplitude displaying no typical resonance-like behavior around the energy region 1.1-1.2 GeV.

[6] The $J/\psi$ and virtual photon from $e^+e^-$ annihilation have negative $C$ parity. The $\kappa'(1150)$ production in the $J/\psi \to \bar{K}^*(892)^0 K^+\pi^-$ decay process is doubly suppressed by $C$ invariance in the $SU(3)_f$ limit and its small $K\pi$ branching ratio. This suppression coincides with the experimental data of quite small production of the $\kappa'(1150)$ compared to the $\kappa(800)$, as is seen in fig.1.

In a future study it is necessary to calculate dynamically the decay $\kappa' \to \kappa(800)\sigma(600) \to K\pi\pi\pi$ in order to obtain a more realistic decay width and also to make the present decay model more effective by examining various strong decay processes.

## References

1. I. Caprini *et al.*, Phys. Rev. Lett. **96**, 132001 (2006); Z.Y. Zhou *et al.*, JHEP **02**, 043 (2005); A. Dobado *et al.*, Phys. Rev. D **56**, 3057 (1997).
2. J.R. Peláez, Mod. Phys. Lett. A **19**, 2879 (2004).
3. S. Descotes-Genon *et al.*, Eur. Phys. J. C **48**, 553 (2006); Z.Y. Zhou *et al.*, Nucl. Phys. A **775**, 212 (2006); H.Q. Zheng *et al.*, Nucl. Phys. A **733**, 235 (2004).
4. E.M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **86**, 770 (2001); E.M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **89**, 121801 (2002).
5. M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **598**, 149 (2004).
6. M. Ablikim *et al.* [BES Collaboration], Phys. Lett. B **633**, 681 (2006).
7. S. Ishida *et al.*, Prog. Theor. Phys. **104**, 785 (2000).
8. S. Ishida *et al.*, Phys. Lett. B **539**, 249 (2002).
9. K. Yamada, in: Proceedings of the Seminar on Perspectives for Studies of Chiral Particles at BES, (eds. W.G. Li *et al.*, IHEP, Beijing, February 2006), 23 (KEK Proceedings 2006-8, November 2006).
10. C. McNeile *et al.* [UKQCD Collaboration], Phys. Rev. D **74**, 014508 (2006).
11. T. Maeda *et al.*, these proceedings.
12. W.-M. Yao *et al.* [Particle Data Group], J. Phys. G **33**, 1 (2006) and 2007 partial update for edition 2008 (URL: http//pdg.lbl.gov).
13. D. Aston *et al.*, Nucl. Phys. **B296**, 493 (1988).
14. H.E. Haber *et al.*, Phys. Rev. D **32**, 2961 (1985).